

# Decreased Cerebrospinal Fluid Allopregnanolone Levels in Women with Posttraumatic Stress Disorder

Ann M. Rasmusson, Graziano Pinna, Prashni Paliwal, David Weisman, Christopher Gottschalk, Dennis Charney, John Krystal, and Alessandro Guidotti

**Background:** Alterations in the  $\gamma$ -amino-butyric acid (GABA) neurotransmitter system have been identified in some populations with posttraumatic stress disorder (PTSD).

**Methods:** To further investigate factors of relevance to GABAergic neurotransmission in PTSD, we measured cerebrospinal fluid (CSF) levels of allopregnanolone and pregnanolone combined (ALLO: congeners that potently and positively modulate effects of GABA at the GABA<sub>A</sub> receptor), 5 $\alpha$ -dihydroprogesterone (5 $\alpha$ -DHP: the immediate precursor for allopregnanolone), dehydroepiandrosterone (DHEA: a negative modulator of GABA<sub>A</sub> receptor function), and progesterone with gas chromatography, mass spectrometry in premenopausal women with ( $n = 9$ ) and without ( $n = 10$ ) PTSD. Subjects were free of psychotropic medications, alcohol, and illicit drugs; all were in the follicular phase of the menstrual cycle except three healthy and four PTSD subjects receiving oral contraceptives.

**Results:** There were no group differences in progesterone, 5 $\alpha$ -DHP, or DHEA levels. The PTSD group ALLO levels were  $< 39\%$  of healthy group levels. The ALLO/DHEA ratio correlated negatively with PTSD re-experiencing symptoms ( $r = -.82$ ,  $p < .008$ ; trend) and with Profile of Mood State depression/dejection scores ( $r = -.70$ ,  $p < .0008$ ).

**Conclusion:** Low CSF ALLO levels in premenopausal women with PTSD might contribute to an imbalance in inhibitory versus excitatory neurotransmission, resulting in increased PTSD re-experiencing and depressive symptoms.

**Key Words:** Posttraumatic stress disorder, depression, allopregnanolone, GABA, dehydroepiandrosterone, women

Progress has been made in treating posttraumatic stress disorder (PTSD) (Aulbacher and Liberzon 2002; Friedman 2002; Resick et al 2002), but the need for improved methods of PTSD prevention and treatment remains and likely will be met only as we expand our understanding of the pathophysiology of PTSD. Research regarding the neurobiology of PTSD thus far has revealed the complexity of the disorder, with reported PTSD-associated alterations in the monoaminergic, corticotropin-releasing factor (CRF), and neuropeptide Y (NPY) transmitter systems (Nutt and Malizia 2004; Southwick et al 2005) as well as in hypothalamic-pituitary-adrenal (HPA) axis (Rasmusson et al 2003; Yehuda et al 2002), thyroid (Karlovic et al 2004; Wang and Mason 1999), and immune system function (Baker et al 2001).

A variety of alterations in the  $\gamma$ -amino-butyric acid (GABA) neurotransmitter system also have been identified in PTSD. Decreased frontal lobe benzodiazepine receptor binding was found in male Viet Nam but not Gulf War veterans with chronic PTSD (Bremner et al 2000a; Fujita et al 2004). Decreased plasma GABA levels were associated with PTSD development in motor vehicle accident victims (Vaiva et al 2004). A polymorphism in the gene for the GABA<sub>A</sub> receptor  $\beta 3$  subunit was associated with higher levels of somatic symptoms, depression, anxiety, and

insomnia in PTSD (Feusner et al 2001). In addition, combat veterans with PTSD (Spivak et al 2000) and refugees from Kosovo who developed PTSD with greater sleep disturbance (Sondergaard et al 2002) showed increased plasma levels of dehydroepiandrosterone (DHEA) and/or its sulfated metabolite (DHEAS), androgenic steroids with negative modulatory effects at brain GABA<sub>A</sub> receptors. Premenopausal women with chronic PTSD also have shown greater DHEA release after maximal adrenal stimulation by adrenocorticotropic hormone (ACTH<sub>1-24</sub>) administration (Rasmusson et al 2004).


In the current study, we further investigated the GABAergic neurotransmitter system in PTSD by measuring, in the cerebrospinal fluid (CSF) of premenopausal women with chronic PTSD, selected neuroactive steroids that influence the expression or function of brain GABA<sub>A</sub> receptors. Two of these steroids, the 3 $\alpha$ -reduced biosynthetic derivatives of progesterone, 3 $\alpha$ -hydroxy-5 $\alpha$ -pregnan-20-one (allopregnanolone) and its stereoisomer, 3 $\alpha$ -hydroxy-5 $\beta$ -pregnan-20-one (pregnanolone) (collectively termed ALLO for this paper), are the most potent and selective positive endogenous modulators of the action of GABA at brain GABA<sub>A</sub> receptors (Lambert et al 2003; Puia et al 1990). In addition, these steroids exert anxiolytic, sedative, and anesthetic effects at nanomolar concentrations (Paul and Purdy 1992), suggesting that they would be logical candidates for a role in the pathophysiology of PTSD.

Methodological challenges have limited previous studies of ALLO in PTSD and other clinical disorders. Radioimmunoassays lack the specificity and sensitivity needed to accurately measure ALLO in the CSF or in the plasma of men, follicular phase premenopausal women, postmenopausal women, and children. Therefore, we used gas-chromatography, mass-spectrometry (GC-MS), a method that allows structural identification and quantification of the neuroactive steroids of interest at femtomolar levels (Uzunova et al 1998). We measured (Figure 1): 1) ALLO; 2) 5 $\alpha$ -dihydroprogesterone (5 $\alpha$ -DHP), the immediate precursor for ALLO that acts with nanomolar potency at progesterone receptors that in turn interact with DNA response elements involved in modifying GABA<sub>A</sub> receptor expression (Canonaco et al 1989; Rupprecht 2003; Rupprecht et al 1993; Schumacher et al

From the Department of Psychiatry (AMR, PP, CG, JK), Department of Neurology (DW, CG), and the Division of Women's Behavioral Health (PP), Yale University School of Medicine, New Haven, Connecticut; Clinical Neuroscience Division (AMR, JK), Veterans Affairs National Center for PTSD, West Haven, Connecticut; Psychiatric Institute (GP, AG), University of Illinois, Chicago, Illinois; and the Departments of Psychiatry, Neuroscience and Pharmacology (DSC), Mount Sinai School of Medicine, New York, New York.

Address reprint requests to Ann M. Rasmusson, MD, VA Connecticut Healthcare System, Psychiatry Service/116A, 950 Campbell Ave., West Haven, CT 06516; E-mail: ann.rasmusson@yale.edu.

Received August 10, 2005; revised March 3, 2006; accepted March 7, 2006.



**Figure 1.** Biosynthetic pathways for brain and/or peripheral neuroactive steroids. HSD, hydroxysteroid dehydrogenase; KSR; ketosteroid reductase; aro, aromatase.

1989); 3) progesterone (PROG), the immediate precursor for 5 $\alpha$ -DHP; and 4) dehydroepiandrosterone (DHEA), a peripherally derived androgen (Compagnone and Mellon 2000) with negative modulatory effects at brain GABA<sub>A</sub> receptors at micromolar concentrations (Park-Chung et al 1999). Dehydroepiandrosterone also has positive modulatory effects at N-methyl-D-aspartate (NMDA) receptors, activity at sigma receptors, and antigluocorticoid as well as neuroprotective properties (Baulieu and Robel 1998).

On the basis of the previous studies' finding of high DHEA or DHEAS levels or reactivity in PTSD and the opposing modulatory effects of DHEA(S) and ALLO at GABA<sub>A</sub> receptors, we hypothesized that CSF ALLO levels would be low and DHEA levels high in women with PTSD. Because the pharmacological properties of these neuroactive steroids might allow them to directly modulate psychiatric symptoms, we also investigated their relationship to clusters of related DSM-IV–defined PTSD symptoms identified by confirmatory factor analysis (Simms et al 2002) (Table 3). We hypothesized that CSF ALLO levels would correlate negatively with the re-experiencing and hyperarousal symptom clusters and that CSF DHEA levels would correlate negatively with the dysphoria symptom cluster (Table 3).

## Methods and Materials

### Subjects

Twenty-seven subjects were recruited for the study via community advertisements. The protocol was approved by the VA Connecticut Healthcare System Human Studies Subcommittee and the Yale Human Investigation Committee; written informed consent was obtained from each subject.

Subjects were screened for current or significant past medical illness and illicit drug use via medical history, physical examination, and laboratory tests. None had identifiable endocrine syndromes, and all reported regular menses on admission to the study.

Subjects were free from psychotropic medications for months to years or had never been treated with such. Subjects abstained from other medications, alcohol, and illicit drugs for at least 4 weeks before lumbar puncture except for four PTSD and three healthy, nontraumatized subjects who used oral contraceptives. One PTSD subject smoked approximately one-half pack of cigarettes per day.

The DSM-IV PTSD Criterion A1/A2 trauma exposure (American Psychiatric Association 1994) was assessed with general and early trauma inventories (ETI) (Bremner et al 2000b) (Table 1). To qualify as a Criterion A2 trauma, physical or sexual trauma ascertained by the ETI had to be scored as "very" or "extremely" distressing and disruptive in at least two of three functional domains (emotional, interpersonal, or school performance).

Healthy, nontraumatized subjects were evaluated psychiatrically with the non-patient edition of the Structured Clinical Interview for DSM-IV (SCID-NP) (First et al 1995a); subjects with a current Axis I diagnosis were excluded. The PTSD subjects were evaluated with the SCID-P (First et al 1995b); subjects with current or past psychotic disorders or bipolar disorder were excluded.

### Menstrual Cycle Monitoring

Menstrual cycle phase was determined by use of a urine test kit that detects the mid-cycle luteinizing hormone surge (Clear Plan Easy, Whitehall Laboratories, Madison, New Jersey) and measurement of plasma progesterone levels 10–14 days later to confirm ovulation. Progesterone levels also were measured in the hospital clinical laboratory at the time of the lumbar puncture to confirm menstrual phase.

### Lumbar Puncture

Subjects fasted except for water intake and abstained from smoking (in the case of the one PTSD subject who smoked) after midnight before the lumbar puncture. Subjects arrived at the Biostudies Unit at 7:30 AM, provided urine for a urine toxicology screen and  $\beta$ -HCG pregnancy testing, and then lay flat. Blood was drawn from an antecubital vein at –30 and –15 min before the lumbar puncture, which was performed between 8:30 and 9:30 AM. The CSF (25 cc) was collected in 1 cc aliquots into polypropylene tubes that were immediately placed on dry ice before storage in a –70° freezer. Blood samples were spun immediately in a refrigerated centrifuge and plasma was stored at –70°C.

### Symptom Ratings

The Profile of Mood States scale (POMS; Educational and Industrial Testing Service, San Diego, California) was rated for the past week by all subjects during the 30 min before lumbar puncture. The Clinician Administered PTSD Scale (CAPS-One Week Version; Blake et al 1993) was administered after the lumbar puncture so as not to influence CSF neuroactive steroid levels by overtly stressing the PTSD subjects with trauma reminders.

### Neuroactive Steroid Measurements

The CSF neuroactive steroid levels were measured with GC-MS. The detection limit for ALLO and the other neurosteroids with this method was approximately 1.0 fmol/mL, and the standard curve was linear between 1 and 10<sup>5</sup> fmol/mL. The intra- and inter-assay coefficients of variation for this method are low (Uzunov et al 1996). Neuroactive steroid measurements were made in CSF aliquot 18 and confirmed either in aliquot 18 or 19. For one subject, CSF was available only from aliquot 19. Average values from multiple measurements of ALLO (either 2 or 3) were used in the data analyses for 15 subjects; only one measurement was available for four subjects. The PROG, 5 $\alpha$ -DHP, and DHEA measurements were made once for each subject.

The CSF samples (700  $\mu$ L) were extracted in ethylacetate and lyophilized; the neurosteroids of interest then were purified and separated with high-performance liquid chromatography (HPLC). Tritiated neurosteroids (New England Nuclear, Boston, Massachusetts) were added to monitor retention time and steroid recovery through the HPLC procedure (Cheney et al 1995), while deuterated internal standards (Cambridge Isotope Laboratories, Andover, Massachusetts) were added to allow quantification of the compound of interest. The HPLC-eluted fractions containing each steroid of

**Table 1.** Demographic Characteristics, PTSD A1, A2 Trauma Exposure, and DSM-IV Psychiatric Diagnoses of Subjects

| Demographic Variable                 | PTSD Subjects<br>(n = 9)   | Healthy Nontraumatized<br>Subjects (n = 10)   | Group Difference            |
|--------------------------------------|--|---|-----------------------------|
| Age                                  | Mean: 33.6 ± 6.1 yrs<br>Range: 25–43 yrs   | Mean: 28.7 ± 8.6 yrs<br>Range: 19–43 yrs  | p < .04 <sup>a</sup>        |
| Weight                               | Mean: 173.3 ± 33.3 lbs   | Mean: 169.7 ± 43.5 lbs  | p > .96                     |
| Education                            | Mean: 14.0 ± 2.0 yrs   | Mean: 15.5 ± 1.8 yrs  | p > .16                     |
| Ethnicity                            | Caucasian: n = 6; 66.7%<br>African American: n = 2; 22.2%<br>Hispanic: n = 1; 11.1%                                  | Caucasian: n = 4; 40%,<br>African American: n = 2; 20%,<br>Hispanic: n = 1; 10%,<br>Asian: n = 2; 20%,<br>Askenazi Jewish: n = 1; 10% |                             |
| Trauma Exposure                      | Age 1st Trauma: All pre-adolescent<br>All with multiple traumas (#2–13)<br>All suffered sexual assault at least once | None  | Yes, by definition.         |
| Psychiatric Diagnoses (No. subjects) |  |   | Yes, by definition.         |
|                                      | Major Depression   | Current/Past<br>4/3   | Bulimia <sup>a</sup><br>0/1 |
|                                      | Dysthymia  | 3/0   |                             |
|                                      | Panic Disorder   | 2/0   |                             |
|                                      | Simple Phobia  | 2/0   |                             |
|                                      | Social Phobia  | 1/0   |                             |
|                                      | Agoraphobia w/o panic  | 2/1   |                             |
|                                      | Generalized Anxiety  | 1/1   |                             |
|                                      | OCD (cleaning only)  | 1/0   |                             |
|                                      | Bulimia  | 0/1   |                             |
|                                      | Anorexia   | 0/1   |                             |
|                                      | Alcohol Dependence <sup>a</sup>  | 0/2   |                             |
|                                      | Marijuana Dependence <sup>a</sup>  | 0/2   |                             |

PTSD, posttraumatic stress disorder; OCD, obsessive compulsive disorder.

<sup>a</sup>> 6 years before study participation (PTSD); > 10 years before study participation (Healthy).

interest then were derivatized for GC-MS (Dong et al 2001; Uzunova et al 1998). Because deuterated DHEA is not available, deuterated allopregnanolone was added to the HPLC-eluted fraction containing DHEA before derivatization. Mass spectrometry analysis was conducted in the standard electron impact mode (EI) for progesterone, ALLO, and DHEA and in the negative ion chemical ionization mode (NICI) with methane as the reaction gas for 5 $\alpha$ -DHP. To calculate the quantity of the neurosteroid of interest in each fraction, the area under the peak of the neurosteroid in the sample was divided by the area under the peak of the deuterated internal standard. For DHEA, the ratio of the area under the DHEA peak to be determined divided by the area under the derivatized ALLO peak was plotted against similar ratios generated in a standard curve prepared by combining known quantities of authentic DHEA (1–200 fmol) with 100 fmol of deuterated ALLO. Further methodological details may be obtained by contacting the authors.

As previously shown, only the GABA<sub>A</sub> receptor active 3 $\alpha$ 5 $\alpha$  or 3 $\alpha$ 5 $\beta$  stereoisomers of ALLO are measurable in CSF, whereas the 3 $\beta$ 5 $\alpha$  and 3 $\beta$ 5 $\beta$  isomers that antagonize the effects of the 3 $\alpha$  isomers at high concentrations are below the limits of detection (Uzunova et al 1998). The 3 $\alpha$ 5 $\alpha$  and 3 $\alpha$ 5 $\beta$  stereoisomers of ALLO were measured together in the current study, because the content of each in CSF is almost identical (Uzunova et al 1998); they are equipotent in modulating GABA action at GABA<sub>A</sub> receptors, and separating them is very time consuming owing to their similar retention times. In the current study, they were eluted with a retention time of approximately 14 min and were easily separated from the 3 $\beta$ 5 $\alpha$  and 3 $\beta$ 5 $\beta$  stereoisomers that have a retention time of approximately 16 minutes.

### Statistical Analyses

The following subjects were excluded from the data analyses: one nontraumatized subject diagnosed with Adjustment Disorder with Depressed Mood in the context of currently stressful family circumstances, one traumatized subject meeting criteria for past and current partial PTSD, a healthy subject from whom CSF could not be obtained during the lumbar puncture, two PTSD and three healthy subjects in the luteal phase of the menstrual cycle at the time of lumbar puncture, and a healthy subject who initially reported regular menses but who failed to ovulate over the next 3 months.

For the purposes of data analysis, subjects receiving oral contraceptives (three healthy and four PTSD subjects) were considered in combination with subjects in the natural follicular phase of the menstrual cycle, because plasma progesterone levels are similar in these states. Analyses were therefore conducted in a total of 9 PTSD and 10 healthy subjects.

Because possible relationships between age, weight, or CSF progesterone levels and CSF levels of the neuroactive steroids of interest have not been defined previously, we performed exploratory Spearman correlations to ascertain whether these factors should be included as covariates in analyses used to compare neuroactive steroid levels between groups.

Nonparametric Mann–Whitney tests were initially used to compare means of the neuroactive steroid levels between groups. Linear regression was used to adjust for age and CSF progesterone levels on the basis of the exploratory analyses. Biologically relevant ratios between CSF levels of selected neuroactive steroids also were computed and compared between the PTSD and healthy groups. The ratio of ALLO to DHEA was calculated as a measure of balance between inhibitory and excitatory influences in the central nervous system (CNS). Ratios between progesterone and 5 $\alpha$ -DHP and between 5 $\alpha$ -DHP and ALLO were calculated to index possible

**Table 2.** Group Differences in CSF Neuroactive Steroid Levels and Ratios

| Findings Relevant to the Pathway for Allopregnanolone Synthesis |                 |       |                     |       |                          |                                     |
|---|-----------------|-------|---------------------|-------|--------------------------|-------------------------------------|
| Neuroactive Steroid Measure                                     | PTSD<br>(n = 9) |       | HEALTHY<br>(n = 10) |       | Unadjusted<br>Comparison | Adjusted <sup>a</sup><br>Comparison |
|   | Mean (fmol/mL)  | SEM   | Mean (fmol/mL)      | SEM   | Mann Whitney <i>p</i>    | Linear Regression <i>p</i>          |
| PROG  | 439.6           | 112.0 | 392.5               | 106.6 | .84                      |                                     |
| 5- $\alpha$ -DHP  | 282.5           | 69.6  | 152.1               | 23.4  | .27                      |                                     |
| ALLO <sup>b</sup>   | 14.1            | 2.6   | 36.2                | 6.6   | .013 <sup>b</sup>        | <i>p</i> < .01 <sup>b</sup>         |
| PROG/5- $\alpha$ -DHP   | 2.91            | .94   | 2.82                | .61   | .780                     |                                     |
| 5- $\alpha$ -DHP/ALLO <sup>b</sup>                              | 28.8            | 9.8   | 5.1                 | .92   | .006 <sup>b</sup>        | <i>p</i> < .04 <sup>b</sup>         |

| Findings Regarding the Balance Between Neuroactive Steroids that Positively and Negatively Modulate GABA <sub>A</sub> Receptor Function |                 |      |                     |      |                          |                                     |
|---|-----------------|------|---------------------|------|--------------------------|-------------------------------------|
| Neuroactive Steroid Measure   | PTSD<br>(n = 9) |      | HEALTHY<br>(n = 10) |      | Unadjusted<br>Comparison | Adjusted <sup>a</sup><br>Comparison |
|   | Mean (fmol/mL)  | SEM  | Mean (fmol/mL)      | SEM  | Mann Whitney <i>p</i>    | Linear Regression <i>p</i>          |
| DHEA  | 220.6           | 34.1 | 212.3               | 46.9 | .60                      |                                     |
| ALLO/DHEA <sup>b</sup>  | .08             | .02  | .22                 | .06  | .008 <sup>b</sup>        | <i>p</i> < .03 <sup>b</sup>         |

PROG, progesterone; DHP, dihydroprogesterone; ALLO, combined allopregnanolone and pregnanolone; DHEA, dehydroepiandrosterone.

<sup>a</sup>Comparison adjusted for age and cerebrospinal fluid (CSF) progesterone levels.

<sup>b</sup>Significant (*p* < .05) difference between posttraumatic stress disorder (PTSD) and Healthy groups.

blocks in the conversion of precursor to product steroid (see Figure 1).

Spearman correlations were performed between CSF ALLO or DHEA levels or the ALLO/DHEA ratio and: 1) clusters of related DSM-IV–defined PTSD symptoms identified by confirmatory factor analysis (Simms et al 2002) in the PTSD subjects, and 2) POMS symptoms scores in all subjects. Bonferroni corrections were used to account for the multiple hypotheses under consideration. It should be noted that Bonferroni corrections are very conservative and pose a risk for Type 2 error especially in this type of situation wherein many of the psychiatric symptoms or neuroactive steroid levels and ratios might have common mechanistic determinants and might be highly correlated.

Descriptive statistics are expressed as group means with SEM.

## Results

### Demographic Characteristics

Subjects' age, weight, education, trauma exposure, and past or current psychiatric diagnoses besides PTSD are tabulated in Table 1. The PTSD and healthy groups differed significantly by age but not by weight or education.

### Ascertainment of Covariates

The CSF progesterone and ALLO levels were positively correlated (all subjects:  $r = .48$ ,  $p < .04$ ; healthy:  $r = .77$ ,  $p < .01$ ; PTSD:  $r = .47$ ,  $p = .20$ ). There was a trend for a negative correlation between age and CSF ALLO levels in all subjects ( $r = -.45$ ,  $p < .06$ ;  $n = 19$ ). This trend occurred in the healthy group considered alone ( $r = -.60$ ;  $p < .07$ ) but not in the PTSD group ( $r = .13$ ,  $p = .73$ ). There were no correlations between body weight and neuroactive steroid levels.

### Differences in Neuroactive Steroid Levels Between PTSD and Healthy Subjects

There were no significant differences between the PTSD and healthy subjects in CSF levels of progesterone, 5- $\alpha$ -DHP, or the progesterone/5- $\alpha$ -DHP ratio (Table 2, Figure 2A). The CSF ALLO levels were significantly lower in the PTSD group (Table 2, Figure 2B), whereas the 5- $\alpha$ -DHP/ALLO ratio was higher. Linear regression

adjusting for age and CSF progesterone levels also revealed group differences in CSF ALLO levels ( $t = -2.77$ ,  $p < .01$ ) and the 5- $\alpha$ -DHP/ALLO ratio ( $t = 2.34$ ,  $p < .04$ ).

The CSF DHEA levels were not different between the PTSD and healthy groups, but the ALLO/DHEA ratio was low in the PTSD subjects (unadjusted comparison:  $p < .008$ ; comparison adjusted for age and CSF progesterone levels:  $t = -2.29$ ,  $p < .03$ ) (Table 2B).

Follesa et al (2002) showed that oral contraceptive administration reduces progesterone and allopregnanolone levels, measured by a less-specific radioimmunoassay method, in the plasma and cerebral cortex of rats and in the plasma of women. In the current study, the PTSD and healthy groups were matched for number of oral contraceptive users. In addition, we performed statistical analyses to determine whether oral contraceptive use might have contributed to the study findings. We found no significant difference in neuroactive steroid levels or their ratios between subjects receiving oral contraception and those in the natural follicular state, when all subjects were considered together or when the PTSD and healthy groups were considered separately. Linear regression analyses controlling for use of oral contraceptives in addition to age and CSF progesterone levels revealed that group differences in CSF ALLO levels ( $p < .04$ ) and the ALLO/DHEA ratio ( $p < .02$ ) persisted. A trend for a group effect on the 5- $\alpha$ -DHP/ALLO ratio was seen in this model ( $p < .07$ ).

### The Relationship Between CSF Neuroactive Steroid Levels and General Mood Symptoms

When all of the subjects were considered together, the CSF ALLO/DHEA ratio correlated strongly and negatively with the POMS depression/dejection and fatigue subscales (Table 3). When the PTSD and healthy subjects were evaluated separately, however, these correlations were preserved in the healthy ( $r = -.76$ ,  $p = .01$  and  $r = -.72$ ,  $p = .02$ , respectively) but not the PTSD group ( $r = .03$ ,  $p = .95$  and  $r = .33$ ,  $p = .38$ , respectively). Nevertheless, when the PTSD group was divided into five subjects with current PTSD alone and four subjects with current comorbid PTSD/MDD, a post hoc analysis revealed that CSF ALLO levels were significantly lower



**Table 3.** Spearman Correlations Between Neuroactive Steroid Levels and Mood/PTSD Symptoms

| Correlations With Profile of Mood State (POMS) Scores in All Subjects (n = 19)                                 |                          |                         |                             |
|--|--------------------------|-------------------------|-----------------------------|
| Profile of Mood State (POMS) Scores  | ALLO                     | DHEA                    | ALLO/DHEA                   |
| Total Poms   | $r = -.45$<br>$p = .05$  | $r = .09$<br>$p = .71$  | $r = -.52$<br>$p = .02$     |
| Anger/Irritation   | $r = -.43$<br>$p = .06$  | $r = .25$<br>$p = .30$  | $r = -.57$<br>$p = .01$     |
| Anxiety/Tension  | $r = -.46$<br>$p = .045$ | $r = .06$<br>$p = .79$  | $r = -.50$<br>$p = .03$     |
| Confusion  | $r = -.31$<br>$p = .20$  | $r = .16$<br>$p = .52$  | $r = -.43$<br>$p = .07$     |
| Depression/Dejection   | $r = -.52$<br>$p = .02$  | $r = .27$<br>$p = .25$  | $r = -.70$<br>$p = .0008^a$ |
| Fatigue  | $r = -.50$<br>$p = .03$  | $r = .30$<br>$p = .21$  | $r = -.63$<br>$p = .004^b$  |
| Vigor  | $r = .23$<br>$p = .34$   | $r = .46$<br>$p = .05$  | $r = .03$<br>$p = .90$      |
| Correlations With PTSD Symptom Clusters in PTSD Subjects (n = 9)   |                          |                         |                             |
| Clusters of Related DSM-IV–Defined PTSD Symptoms Identified by Confirmatory Factor Analysis (Simms et al 2002) | ALLO                     | DHEA                    | ALLO/DHEA                   |
| Re-Experiencing: Symptoms in this cluster are the same as DSM-IV Criterion B PTSD re-experiencing symptoms     | $r = -.72$<br>$p = .03$  | $r = .55$<br>$p = .13$  | $r = -.82$<br>$p = .007^c$  |
| B1-intrusive thoughts  |                          |                         |                             |
| B2-nightmares  |                          |                         |                             |
| B3-flashbacks  |                          |                         |                             |
| B4-emotional distress at trauma cue exposure   |                          |                         |                             |
| B5-physiological reactivity to trauma cues   |                          |                         |                             |
| Active Avoidance   | $r = -.08$<br>$p = .83$  | $r = .23$<br>$p = .56$  | $r = -.24$<br>$p = .53$     |
| C1-avoidance of trauma-related thoughts, feelings, conversions   |                          |                         |                             |
| C2-avoidance of people, places, things   |                          |                         |                             |
| Dysphoria  | $r = .03$<br>$p = .95$   | $r = -.18$<br>$p = .65$ | $r = .04$<br>$p = .91$      |
| C3-amnesia   |                          |                         |                             |
| C4-decreased interest in everyday activities   |                          |                         |                             |
| C5-detachment or estrangement from others  |                          |                         |                             |
| C6-restricted range of affect  |                          |                         |                             |
| C7-sense of foreshortened future   |                          |                         |                             |
| D1-disturbed sleep   |                          |                         |                             |
| D2-irritability  |                          |                         |                             |
| D3-poor concentration  |                          |                         |                             |
| Arousal  | $r = .78$<br>$p = .01$   | $r = -.21$<br>$p = .59$ | $r = .65$<br>$p = .06$      |
| D4-hypervigilance  |                          |                         |                             |
| D5-increased startle   |                          |                         |                             |

Abbreviations as in Table 2.

<sup>a</sup>A significant ( $p < .0024$ ) correlation after Bonferroni correction for 21 comparisons.

<sup>b</sup>Trend for a significant ( $.0024 < p < .0048$ ) correlation after Bonferroni correction for 21 comparisons.

<sup>c</sup>A trend ( $p < .008$ ) for a significant correlation after Bonferroni correction for 12 comparisons.

in the PTSD/current MDD group ( $7.7 \pm 4.6$  fmol/mL vs.  $19.3 \pm 5.4$  fmol/mL, Mann–Whitney  $p = .015$ ).


### The Relationship Between CSF Neuroactive Steroid Levels and PTSD Symptoms

In the small sample of nine PTSD subjects, there was a trend for a strong negative Spearman correlation between the CSF ALLO/DHEA ratio and PTSD re-experiencing symptoms ( $r = -.82$ ,  $p < .008$ , Table 3 and Figure 3). In addition, a post hoc analysis revealed that re-experiencing symptoms were significantly higher in the PTSD/current MDD group compared with the PTSD alone group ( $16.2 \pm 2.2$  vs.  $8.6 \pm 5.5$  fmol/mL, respectively; Mann–Whitney test  $p = .04$ ). As noted in the previous section, the PTSD/current MDD group also had lower CSF ALLO levels.

## Discussion

### Summary of Findings

Women with PTSD who were free of psychotropic medications for several months or longer were found to have decreased CSF ALLO levels that were only 39% of healthy group levels, whereas CSF DHEA levels were normal. If levels of these neuroactive steroids in CSF reflect levels in brain, the data suggest that the balance between CNS inhibitory and excitatory neurosteroids might be diminished in the women with PTSD. Consistent with this possibility was a strong negative correlation between the CSF ALLO/DHEA ratio (but not ALLO levels alone in this small sample) and depressive symptoms among all subjects as well as a trend for a strong negative relationship between the ALLO/DHEA ratio and




**Figure 2.** (A) Relative cerebrospinal fluid levels of progesterone (PROG), 5α-dihydroprogesterone (5α-DHP), and 3α5α and 3α5β-tetrahydroprogesterone (ALLO) in 9 women with posttraumatic stress disorder (PTSD) and 10 healthy nontraumatized women; (B) a scattergram of the CSF ALLO levels in the 10 healthy subjects (black circles) and 9 PTSD subjects (red squares).


PTSD re-experiencing symptoms among the PTSD subjects. In addition, CSF ALLO levels were lower and PTSD re-experiencing symptoms higher in the PTSD subjects with current comorbid MDD compared with the PTSD subjects without current MDD. This study thus extends previous work finding low CSF ALLO levels in male and female patients with MDD (Uzunova et al 1998) to a cohort of premenopausal women with PTSD.

**Mechanisms That Might Produce Low ALLO Levels in Women With PTSD**

In the only work thus far to model reductions in brain ALLO levels in response to chronic stress, Dong et al (2001) found decreased brain ALLO levels in male mice exposed to prolonged social isolation, an effect reproduced in testosterone-treated ovariectomized female mice but not in normal female mice (Pinna et al 2005). In this model, decreased brain ALLO levels resulted from downregulation of 5α-reductase gene expression and inadequate



**Figure 3.** Relationship between posttraumatic stress disorder (PTSD) re-experiencing symptoms and the ratio of cerebrospinal fluid (CSF) 3α5α and 3α5β-tetrahydroprogesterone (ALLO) levels to CSF dehydroepiandrosterone (DHEA) levels in nine women with PTSD (Spearman  $r = -.82$ ,  $p < .008$ ; trend with Bonferroni corrections).



**Figure 4.** The ratio of 5α-dihydroprogesterone (5α-DHP) to 3α,5α and 3α,5β-tetrahydroprogesterone (ALLO) in the cerebrospinal fluid of individual posttraumatic stress disorder and healthy, nontraumatized premenopausal female subjects.

provision of the precursor steroid, 5α-DHP, for conversion into ALLO (see Figure 1).

The decrease in CSF ALLO levels in the women with PTSD examined in the current study cannot, however, be attributed to a downregulation of 5α-reductase levels or function. Progesterone and 5α-DHP levels as well as the ratio between progesterone and 5α-DHP were not different between the PTSD and healthy groups (Table 2 and Figure 2). Instead, the 5α-DHP/ALLO ratio was high in the women with PTSD (Figure 4). Thus, we believe that the low ALLO levels in the women with PTSD are due to either an enhancement of ALLO conversion to unmeasured inactive metabolites or a reduced conversion of 5α-DHP to ALLO. Previous work by Uzunova et al (1998) showed that levels of 3α,5α,20α-hexahydroprogesterone, one of the inactive metabolites of allopregnanolone, were below the limits of detection (< 1 fmol/mL) in the CSF of healthy and depressed subjects. Although it is possible that this metabolite could be increased in the CSF of our PTSD sample (we did not measure it), we believe that reduced conversion of 5α-DHP to ALLO is the most likely explanation for our findings.

There are several mechanisms by which conversion of 5α-DHP to ALLO could be deficient in PTSD. A combination of these mechanisms might operate within an individual, or different mechanisms might operate within different individuals. Because the enzyme that converts 5α-DHP to ALLO, 3α-hydroxysteroid dehydrogenase (3α-HSD), operates in a bidirectional manner, it is possible that faulty inhibition of 3α-HSD oxidase activity by accumulated reduced nicotinamide adenine dinucleotide phosphate (NADPH) under conditions of stress could result in preferential conversion of ALLO to 5α-DHP and deficient ALLO levels during stress (Penning et al 2004). Alternatively, a functional polymorphism or frank mutation of the 3α-HSD gene or dysregulation of 3α-HSD gene expression could lead to reduced CSF ALLO levels (Hou et al 1998).

### Properties of Allopregnanolone of Possible Relevance to PTSD

In the CNS, ALLO acts in the nanomolar range as a positive allosteric modulator of GABA action at GABA<sub>A</sub> receptors (Lambert et al 2003; Puia et al 1990); ALLO also inhibits nicotinic acetylcholine receptor-mediated currents, although the behaviorally inert 3 $\beta$  isomer of ALLO is equipotent in this respect. Thus the anxiolytic, anticonflict, anticonvulsant, anesthetic, sedative, and neuroprotective actions of ALLO (Ciriza et al 2004; Djebaili et al 2005; Kaminski et al 2003; Rhodes et al 2004) are all believed to be a consequence of ALLO's action at GABA<sub>A</sub> receptors (Guidotti et al 2001; Pinna et al 2000; Puia et al 2003).

Recent work suggests that although neurosteroids such as ALLO have activity at all subtypes of GABA<sub>A</sub> receptors, they have highest affinity for a benzodiazepine-resistant subset of extrasynaptic GABA<sub>A</sub> receptors composed of  $\alpha_4$  and  $\delta$  subunit combinations or  $\alpha_6$ ,  $\gamma$ , and  $\beta$  subunit combinations (Lambert et al 2003). These extrasynaptic receptors are activated by concentrations of GABA lower than that required for activation of synaptically located GABA<sub>A</sub> receptors. As a consequence, they are thought to maintain a tonic inhibitory conductance that modulates gain in neuronal output during periods of increased input (Mitchell and Silver 2003; Mody and Pearce 2004; Semyanov et al 2003, 2004). Of note,  $\alpha_4$ ,  $\delta$ , and  $\alpha_6$  GABA<sub>A</sub> receptor subunits increase under conditions in which allopregnanolone levels are decreased (Follesa et al 2001; Gulinello et al 2002; Pinna et al 2004; Smith et al 1998; Sundstrom-Poromaa et al 2002). These putative “compensatory” receptor changes might be inadequate, however, because anxious behavior, irritability, and aggression are observed to increase under these circumstances.

We thus speculate that an uncompensated defect in the synthesis or disposition of ALLO with resultant reductions in tonic GABA-mediated inhibitory conductance might contribute to the pathophysiology of PTSD, at least in some individuals.

### Low ALLO Levels and PTSD Neurobiological Phenotypes

Low CSF ALLO levels or a low ratio of ALLO to DHEA, resulting in a relative decrease in CNS inhibitory versus excitatory tone, could contribute directly to hyperreactivity of the amygdala (Ressler et al 2002; Sun et al 2004) as well as increased extrahypothalamic CRF (Patchev et al 1994) and monoaminergic responses (Goldstein et al 1996; Southwick et al 2005) during trauma exposure or exposure to trauma reminders. A decrease in extrasynaptic GABA<sub>A</sub> receptor-mediated tonic inhibition also might facilitate phasic increases in glutamatergic and GABAergic activity that could, for instance, disrupt frontal lobe-mediated inhibition of the amygdala. This might contribute to an enhancement of fear conditioning as well as resistance to extinction of fearful conditioned responses and promote the development of PTSD (Rosenkranz et al 2003; Southwick et al 2005). Because allopregnanolone is thought to provide neuroprotection, subjects with a diminished capacity for ALLO release also might be prone to stress-induced structural and functional damage of brain areas such as the hippocampus—as seen in some populations with PTSD (Bonne et al 2001; De Bellis et al 2001; Gilbertson et al 2002; Hull 2002).

Dysregulation of the HPA axis in PTSD (Rasmusson et al 2003; reviewed in Yehuda et al 2002; Young and Breslau 2004) also might be related to low CNS ALLO levels. Decreased cortisol release during stress could interfere with upregulation of the 3 $\alpha$ -HSD gene and limit adaptive increases in ALLO production during stress (Hou et al 1997). On the other hand, ALLO release during stress provides delayed homeostatic control over activation of the HPA axis (Barbaccia et al 2001; Patchev et al 1996) and reduces expression of the

CRF and arginine vasopressin (AVP) genes in the hypothalamus (Patchev et al 1994, 1996). Thus, low ALLO levels might account for the increases in HPA axis reactivity and increases in 24-hour urinary cortisol output observed in some populations with PTSD, such as premenopausal women with comorbid lifetime or current MDD (Heim et al 2000; Lipschitz et al 2003; Rasmusson et al 2001; Young and Breslau 2004).

### Study Limitations

Although the current study is methodologically rigorous and the findings are intrinsically robust and appealing from a theoretical point of view, the small sample size constitutes a significant limitation by increasing the risk for Type II error and limiting the confidence with which the findings might be generalized. The relationship between age and CSF ALLO levels and possible effects of oral contraceptives on CSF ALLO levels (Follesa et al 2002) should be investigated again in larger human populations. In addition, smoking effects on ALLO levels should be examined. The only subject in the study who smoked had the highest ALLO level in the PTSD group. Because high-dose nicotine increases brain ALLO levels in the rat (Porcu et al 2003), it is possible that heavy smoking reduces PTSD symptoms via effects on ALLO levels, accounting in part for the increased rates of high-intensity smoking in PTSD.

The current study also did not measure all neuroactive steroids that modulate GABA<sub>A</sub> receptor function. In earlier work, we attempted to measure CSF allotetrahydrodeoxy-corticosterone (THDOC: a potent positive modulator of GABA at GABA<sub>A</sub> receptors) and 3 $\alpha$ -diol (a weak androgenic metabolite of the potent androgen, 5 $\alpha$ -dihydrotestosterone, and a potent positive modulator of GABA at GABA<sub>A</sub> receptors), both presumably of peripheral origin. Both were below the limits of detection. Other negative modulators of GABA<sub>A</sub> receptor function such as DHEAS, which acts more potently than DHEA at both NMDA and GABA<sub>A</sub> receptors, and pregnenolone sulfate (Baulieu and Robel 1998; Park-Chung et al 1999) can and should be assessed in future studies.

We also cannot tell from the current study whether the CSF neurosteroid levels measured constitute baseline or stress levels induced by undergoing lumbar puncture or to what extent brain, spinal cord (Patte-Mensah et al 2004), or peripheral sources might have contributed to them. Previous work suggests that the adrenal gland contributes significantly to stress-induced increases in brain and, presumably, CSF ALLO levels (Barbaccia et al 1996; Purdy et al 1991; Vallee et al 2000). This assertion, however, is based on the capacity of adrenalectomy to reduce stress-related increases in brain allopregnanolone levels and does not take into account the possibility that adrenalectomy-related corticosterone reductions might disrupt DNA glucocorticoid response element-mediated maintenance of brain 3 $\alpha$ -HSD gene transcription (Hou et al 1997). Be that as it may, future studies using continuous CSF sampling or maneuvers that markedly increase plasma levels of adrenally derived steroids before performance of lumbar puncture might help us learn to what extent peripheral neuroactive steroids contribute to CSF steroid levels in humans.

The high degree of comorbidity between PTSD and current MDD observed in the study is consistent with rates of PTSD/MDD comorbidity observed in epidemiological studies (e.g., Kessler et al 1995). This makes it difficult to know whether the low CSF ALLO levels observed in the PTSD subjects were attributable to PTSD or depression. In fact, our data suggest that CSF ALLO levels might vary continuously with depressive symptoms. Not only did we observe a strong negative correlation between depressive symptoms and the ALLO/DHEA ratio among all subjects, but CSF ALLO levels were highest in healthy subjects having no history of MDD, lower in

PTSD subjects with past but not current MDD, and lowest in PTSD subjects with current MDD. In contrast, we observed a relationship between low CSF ALLO levels and PTSD re-experiencing symptoms, only one of which is a symptom of MDD. Second, PTSD/MDD typically constitutes more severe PTSD, owing to the fact that four PTSD symptoms are also symptoms of MDD. Third, whereas a previous diagnosis of depression has been shown to predispose to PTSD, depression diagnosed for the first time after trauma exposure is almost always associated with PTSD (Breslau et al 2000). In addition, it is unclear whether subjects in previous studies of ALLO levels in MDD were assessed for trauma exposure or comorbid PTSD (Romeo et al 1998; Uzunova et al 1998).

Finally, as previously noted, animal studies have shown that brain ALLO levels increase in response to acute stress but decrease in response to the stress of chronic social isolation. The cross-sectional design of the current study, however, did not allow us to discriminate whether low ALLO levels were pre-existing (perhaps associated with MDD) and predisposed to the development of PTSD after trauma, resulted from traumatic stress exposure, or emerged as a consequence of social isolation experienced in the aftermath of trauma—possibilities that are not mutually exclusive. Prospective studies will be necessary to evaluate these possibilities. Future studies also will be necessary to determine whether low ALLO levels are diagnosis-specific or influence patterns of symptoms or severity of illness across psychiatric disorders.

### Significance

The prevalence of PTSD is about 8% in the general population (Kessler et al 1995) but reaches above 15% in combat veterans (Hoge et al 2004; Kessler et al 1995; Kulka et al 1990), victims of rape and violent assault (Breslau et al 1998; Deykin and Buka 1997; Kessler et al 1995), and adolescents exposed to compound community trauma (Fitzpatrick and Boldizar 1993; Horowitz et al 1995; Lipschitz et al 2000). In addition, the prevalence of PTSD is at least twice as high among adult women compared with men when trauma exposure is controlled (Breslau et al 1998; Deykin and Buka 1997; Giaconia et al 1995). Work that advances our understanding of factors that mediate decreases in ALLO in premenopausal women with PTSD might spur the development of new methods for the prevention or targeted treatment of this prevalent, disabling, and costly disorder as well as its comorbid conditions.

For example, women with PTSD/MDD are less likely than women with PTSD alone to recover fully in response to either prolonged exposure or cognitive processing therapy (Nishith et al 2005). Perhaps raising ALLO levels in these patients would increase treatment responsiveness. For instance, serotonin selective reuptake inhibitors (SSRIs), one class of medications effective in some individuals with PTSD (Davidson et al 2004; Stein et al 2000), have been shown to increase brain levels of ALLO in animals (Pinna et al 2003) and CSF and plasma ALLO levels in humans (Griffin and Mellon 1999; Romeo et al 1998; Trauger et al 2002; Uzunova et al 1998). Interestingly, two PTSD subjects from the current study were SSRI nonresponders. These subjects had the second and third highest 5 $\alpha$ -DHP/ALLO ratios, suggesting that more extreme blocks in the conversion of 5 $\alpha$ -DHP to ALLO might confer SSRI treatment resistance. Perhaps the administration of synthetic ALLO-like compounds currently under development (e.g., Gulinello et al 2003; Kaminski et al 2004) or medications such as topiramate that act, in part, directly at extrasynaptic GABA<sub>A</sub> receptors (Berlant 2004; Johnson et al 2005) would benefit such individuals. The use of other medications that increase plasma ALLO levels, perhaps by other mechanisms, also might be found to be effective in PTSD or other

psychiatric disorders on this basis (Genazzani et al 2003; Marx et al 2003; Schmidt et al 2005; Strous et al 2003; Wolkowitz et al 1999).

### Conclusions

Future work aimed at elucidating the mechanisms responsible for low ALLO levels in women with PTSD and ascertaining the extent to which deficits in ALLO production generalize to other PTSD populations might lead to the development of new and more effective interventions for the prevention and treatment of PTSD and its comorbid disorders.

*This work was supported by the Veterans Affairs (VA) National Center for PTSD, Clinical Neurosciences Division in West Haven Connecticut, the Yale Women's Behavioral Health Program, a VA Merit Review (to AMR), National Institute of Mental Health grants MH 49486 and 56890 (to AG), and National Institute on Drug Abuse number 1K12DA14038-01 grant support (to AMR).*

*We thank Valinda Fox, RN, Barbara Corn, PhD., Willie Ford, Angelina Genovese, RNC, M.B.A., Elizabeth O'Donnell, RN, and Brenda Breault, RN, M.A., for their professional and technical support.*

- Albucher RC, Liberzon I (2002): Psychopharmacological treatment in PTSD: A critical review. *J Psychiatric Res* 36:355–367.
- American Psychiatric Association (1994): *Diagnostic and Statistical Manual of Mental Disorders*, 4<sup>th</sup> ed. Washington, DC: American Psychiatric Association.
- Baker DG, Ekhaton NN, Kasckow JW, Hill KK, Zoumakis E, Dashevsky BA, et al (2001): Plasma and cerebrospinal fluid interleukin-6 concentrations in posttraumatic stress disorder. *Neuroimmunomodulation* 9:209–217.
- Barbaccia ML, Serra M, Purdy RH, Biggio G (2001): Stress and neuroactive steroids. *Int Rev Neurobiol* 46:243–272.
- Baulieu E-E, Robel P (1998): Dehydroepiandrosterone (DHEA) and dehydroepiandrosterone sulfate (DHEAS) as neuroactive neurosteroids. *Proc Natl Acad Sci U S A* 95:4089–4091.
- Berlant JL (2004): Prospective open-label study of add-on and monotherapy topiramate in civilians with chronic nonhallucinatory posttraumatic stress disorder. *BMC Psychiatry* 4:24.
- Blake DD, Weathers FW, Nagy LM (1993): A clinician rating scale for assessing current and lifetime PTSD: The CAPS-1. *Behav Ther* 18:187–188.
- Bonne O, Brandes D, Gilboa A, Gomori JM, Shenton ME, Pitman RK, Shalev AY (2001): Longitudinal MRI study of hippocampal volume in trauma survivors with PTSD. *Am J Psychiatry* 158:1248–1251.
- Bremner JD, Innis RB, Southwick SM, Staib L, Zoghbi S, Charney DS (2000a): Decreased benzodiazepine receptor binding in prefrontal cortex in combat-related posttraumatic stress disorder. *Am J Psychiatry* 157: 1120–1126.
- Bremner JD, Vermetten E, Mazure CM (2000b): Development and preliminary psychometric properties of an instrument for the measurement of childhood trauma: The Early Trauma Inventory. *Depress Anxiety* 12:1–12.
- Breslau N, Davis GC, Peterson EL, Schultz LR (2000): A second look at comorbidity in victims of trauma: The posttraumatic stress disorder-major depression connection. *Biol Psychiatry* 48:902–909.
- Breslau N, Kessler RC, Chilcoat HD, Schultz LR, Davis GC, Andreski P (1998): Trauma and posttraumatic stress disorder in the community: The 1996 Detroit area survey of trauma. *Arch Gen Psychiatry* 55:626–632.
- Canonaco M, O'Connor LH, Pfaff DW, McEwen BS (1989): Longer term progesterone treatment induces changes of GABAA receptor levels in forebrain sites in the female hamster: Quantitative autoradiography study. *Exp Brain Res* 77:407–411.
- Cheney DL, Uzunov D, Costa E, Guidotti A (1995): Gas chromatographic-mass fragmentographic quantitation of 3 alpha-hydroxy-5 alpha-pregnan-20-one (allopregnanolone) and its precursors in blood and brain of adrenalectomized and castrated rats. *J Neurosci* 15:4641–4650.
- Ciriza I, Azcoitia I, Garcia-Segura LM (2004): Reduced progesterone metabolites protect rat hippocampal neurons from kainic acid excitotoxicity in vivo. *J Neuroendocrin* 16:58–63.
- Compagnone NA, Mellon SH (2000): Neurosteroids: Biosynthesis and function of these novel neuromodulators. *Front Neuroendocrinol* 21:1–56.



- Davidson J, Landerman LR, Clary CM (2004): Improvement of anger at one week predicts the effects of sertraline and placebo in PTSD. *J Psychiatric Res* 38:497–502.
- De Bellis MD, Hall J, Boring AM, Frustaci K, Moritz G (2001): A pilot longitudinal study of hippocampal volumes in pediatric maltreatment-related posttraumatic stress disorder. *Biol Psychiatry* 50:305–309.
- Daykin EY, Buka SL (1997): Prevalence and risk factors for posttraumatic stress disorder among chemically dependent adolescents. *Am J Psychiatry* 154:752–757.
- Djebaili M, Guo Q, Pettus EH, Hoffman SW, Stein DG (2005): The neurosteroids progesterone and allopregnanolone reduce cell death, gliosis, and functional deficits after traumatic brain injury in rats. *J Neurotrauma* 22:106–118.
- Dong E, Matsumoto K, Uzunova V, Sugaya I, Takahata H, Watanabe H, et al (2001): Brain 5 $\alpha$ -dihydroprogesterone and allopregnanolone synthesis in a mouse model of protracted social isolation. *Proc Natl Acad Sci U S A* 98:2849–2854.
- Feusner J, Ritchie T, Lawford B, Young RM, Kann B, Noble EP (2001): GABA(A) receptor beta 3 subunit gene and psychiatric morbidity in a post-traumatic stress disorder population. *Psychiatry Res* 104:109–117.
- First MB, Spitzer RL, Gibbon M, Williams JB (1995a): *Structured Clinical Interview for the DSM-IV Axis I Disorders-NonPatient Edition (SCID-I/NP, Version 2.0)*. New York: Biometrics Research Dept., New York State Psychiatric Institute.
- First MB, Spitzer RL, Gibbon M, Williams JB (1995b): *Structured Clinical Interview for the DSM-IV Axis I Disorders-Patient Edition (SCID-I/P, Version 2.0)*. New York: Biometrics Research Dept., New York State Psychiatric Institute.
- Fitzpatrick KM, Boldizar JP (1993): The prevalence and consequences of exposure to violence among African-American youths. *J Am Acad Child Adol Psychiatry* 32:424–430.
- Follesa P, Concas A, Porcu P, Sanna E, Serra M, Mostallino MC, et al (2001): Role of allopregnanolone in regulation of GABA(A) receptor plasticity during long-term exposure to and withdrawal from progesterone. *Brain Res Brain Res Rev* 37:81–90.
- Follesa P, Porcu P, Sogliano C, Cinus M, Biggio F, Mancuso L, et al (2002): Changes in GABA $\alpha$  receptor gamma 2 subunit gene expression induced by long-term administration of oral contraceptives in rats. *Neuropharm* 42:325–336.
- Friedman MJ (2002): Future pharmacotherapy for post-traumatic stress disorder: Prevention and treatment. *Psychiatric Clin N Am* 25:427–441.
- Fujita M, Southwick SM, Denucci CC, Zoghbi SS, Dillon MS, Baldwin RM, et al (2004): Central type benzodiazepine receptors in Gulf War veterans with posttraumatic stress disorder. *Biol Psychiatry* 56:95–100.
- Genazzani AD, Stomati M, Bernardi F, Pieri M, Rovati L, Genazzani AR (2003): Long-term low-dose dehydroepiandrosterone oral supplementation in early and late postmenopausal women modulates endocrine parameters and synthesis of neuroactive steroids. *Fertil Steril* 80:1495–1501.
- Giaconia RM, Reinherz HZ, Silverman AB, Pakiz B, Frost AK, Cohen E (1995): Traumas and posttraumatic stress disorder in a community population of older adolescents. *J Am Acad Child Adol Psychiatry* 34:1369–1379.
- Gilbertson MW, Shenton ME, Ciszewski A, Kasai K, Lasko NB, Orr SP, Pitman RK (2002): Smaller hippocampal volume predicts pathologic vulnerability to psychological trauma. *Nat Neurosci* 5:1242–1247.
- Goldstein LE, Rasmusson AM, Bunney BS, Roth RH (1996): Role of the amygdala in the coordination of behavioral, neuroendocrine, and prefrontal cortical monoamine responses to psychological stress in the rat. *J Neurosci* 16:4787–98.
- Griffin LD, Mellon SH (1999): Selective serotonin reuptake inhibitors directly alter activity of neurosteroidogenic enzymes. *Proc Natl Acad Sci U S A* 96:13512–13517.
- Guidotti A, Dong E, Matsumoto K, Pinna G, Rasmusson AM, Costa E (2001): The socially-isolated mouse: A model to study the putative role of allopregnanolone and 5 $\alpha$ -dihydroprogesterone in psychiatric disorders. *Brain Res Rev* 37:110–115.
- Gulinello M, Gong QH, Smith SS (2002): Progesterone withdrawal increases the alpha4 subunit of the GABA(A) receptor in male rats in association with anxiety and altered pharmacology – a comparison with female rats. *Neuropharmacology* 43:701–714.
- Gulinello M, Gong QH, Smith SS (2003): Progesterone withdrawal increases the anxiolytic actions of gaboxadol: Role of alpha4betadelta GABA(A) receptors. *Neuroreport* 14:43–46.
- Heim C, Newport DJ, Heit S, Graham YP, Wilcox M, Bonsall R, et al (2000): Pituitary-adrenal and autonomic responses to stress in women after sexual and physical abuse in childhood. *JAMA* 284:592–597.
- Hoge CW, Castro CA, Messer SC, McGurk D, Cotting DI, Koffman RL (2004): Combat duty in Iraq and Afghanistan, mental health problems, and barriers to care. *New Engl J Med* 351:13–22.
- Horowitz K, Weine S, Jekel J (1995): PTSD symptoms in urban adolescent girls: Compounded community trauma. *J Am Child Adol Psychiatry* 34:1353–1361.
- Hou YT, Lin HK, Penning TM (1998): Dexamethasone regulation of the rat 3-hydroxysteroid/dihydrodiol dihydrogenase (3-HSD/DD) gene. *Mol Pharmacol* 53:459–466.
- Hull AM (2002): Neuroimaging findings in post-traumatic stress disorder. Systematic review. *Br J Psychiatry* 181:102–110.
- Johnson BA, Ait-Daoud N, Akhtar FZ, Javors MA (2005): Use of oral topiramate to promote smoking abstinence among alcohol-dependent smokers: A randomized controlled trial. *Arch Int Med* 165:1600–1605.
- Kaminski RM, Gasior M, Carter RB, Witkin JM (2003): Protective efficacy of neuroactive steroids against cocaine kindled-seizures in mice. *Eur J Pharm* 474:217–222.
- Kaminski RM, Livingood MR, Rogawski MA (2004): Allopregnanolone analogs that positively modulate GABA receptors protect against partial seizures induced by 6-Hz electrical stimulation in mice. *Epilepsia* 45:864–867.
- Karlovic D, Marusic S, Martinac M (2004): Increase of serum triiodothyronine concentration in soldiers with combat-related chronic post-traumatic stress disorder with or without alcohol dependence. *Wiener Klinische Wochenschrift* 116:385–390.
- Kessler RC, Sonnega A, Bromet E, Hughes M, Nelson CB (1995): Posttraumatic stress disorder in the National Comorbidity Survey. *Arch Gen Psychiatry* 52:1048–1060.
- Koenen KC, Hitsman B, Lyons MJ, Niaura R, McCaffery J, Goldberg J, et al (2005): A twin registry study of the relationship between posttraumatic stress disorder and nicotine dependence in men. *Arch Gen Psychiatry* 62:1258–1265.
- Kulka RA, Schlenger WE, Fairbank JA, Hough RL, Jordan BK, Marmar CR, Weiss DS (1990): *Trauma and the Vietnam War Generation*. New York: Brunner/Mazel.
- Lambert JJ, Bellelli D, Peden DR, Vardy AW, Peters JA (2003): Neurosteroid modulation of GABA $\alpha$  receptors. *Prog Neurobiol* 71:67–80.
- Lipschitz DS, Rasmusson AM, Yehuda R, Wang S, Anyan W, Gueogueieva R, Grilo C, Fehon DC, Southwick SM (2003): Salivary cortisol responses to dexamethasone in adolescents with posttraumatic stress disorder. *J Am Acad Child Adol Psychiatry* 42:1310–1317.
- Lipschitz DS, Rasmusson AM, Anyan W, Cromwell P, Southwick SM (2000): Clinical and functional correlates of posttraumatic stress disorder in urban adolescent girls in a primary care clinic. *J Am Child Adol Psychiatry* 39:1104–1111.
- Marx CE, VanDoren MJ, Duncan GE, Lieberman JA, Morrow AL (2003): Olanzapine and clozapine increase the GABAergic neuroactive steroid allopregnanolone in rodents. *Neuropsychopharm* 28:1–13.
- Mitchell SJ, Silver RA (2003): Shunting inhibition modulates neuronal gain during synaptic excitation. *Neuron* 38:433–445.
- Mody I, Pearce RA (2004): Diversity of inhibitory neurotransmission through GABA $\alpha$  receptors. *TINS* 27:569–575.
- Nishith P, Nixon RDV, Resick PA (200): Resolution of trauma-related guilt following treatment of PTSD in female rape victims: A result of cognitive processing therapy targeting comorbid depression? *J Affective Dis* 86:259–86265.
- Nutt DJ, Malizia AL (2004): Structural and functional brain changes in post-traumatic stress disorder. *J Clin Psychiatry* 65(suppl 1):11–17.
- Park-Chung M, Malayev A, Purdy RH, Gibbs TT, Farb D (1999): Sulfated and unsulfated steroids modulate g-aminobutyric acidA receptor function through distinct sites. *Brain Res* 830:72–87.
- Patchev VK, Hassan AH, Holsboer DF, Almeida OF (1996): The neurosteroid tetrahydroprogesterone attenuates the endocrine response to stress and exerts glucocorticoid-like effects on vasopressin gene transcription in the rat hypothalamus. *Neuropsychopharm* 15:533–540.
- Patchev VK, Shoaib M, Holsboer F, Almeida OF (1994): The neurosteroid tetrahydroprogesterone counteracts corticotropin-releasing hormone-induced anxiety and alters the release and gene expression of corticotropin-releasing hormone in the rat hypothalamus. *Neurosci* 62:265–271.

- Patte-Mensah C, Penning TM, Mensah-Nyagan AG (2004): Anatomical and cellular localization of neuroactive 5 $\beta$ -reduced steroid-synthesizing enzymes in the spinal cord. *J Comp Neurol* 477:286–299.
- Paul SM, Purdy RH (1992): Neuroactive steroids. *FASEB J* 6:2311–2322.
- Penning TM, Jin Y, Steckelbroeck S, Lanisnik Rizner T, Lewis M (2004): Structure–function of human 3-hydroxysteroid dehydrogenases genes and proteins. *Mol Cell Endocrin* 215:63–72.
- Pinna G, Agis-Balboa RC, Doueiri M-S, Guidotti A, Costa E (2004): Brain neurosteroids in gender-related aggression induced by social isolation. *Crit Rev Neurobiol* 16:75–82.
- Pinna G, Costa E, Guidotti A (2005): Changes in brain testosterone and allopregnanolone biosynthesis elicit aggressive behavior. *Proc Natl Acad Sci U S A* 102:2135–2140.
- Pinna G, Dong E, Matsumoto K, Costa E, Guidotti A (2003): In socially isolated mice, the reversal of brain allopregnanolone down-regulation mediates the anti-aggressive action of fluoxetine. *Proc Natl Acad Sci U S A* 100:2035–2040.
- Pinna G, Uzunova V, Matsumoto K, Puia G, Mienville JM, Costa E, Guidotti A (2000): Brain allopregnanolone regulates the potency of the GABA(A) receptor agonist muscimol. *Neuropharm* 39:440–448.
- Porcu P, Sogliano C, Cinus M, Purdy RH, Biggio G, Concas A (2003): Nicotine-induced changes in cerebrocortical neuroactive steroids and plasma corticosterone concentrations in the rat. *Pharmacol Biochem Behav* 74:683–690.
- Puia G, Mienville J-M, Matsumoto K, Takahata H, Watanabe H, Costa E, Guidotti A (2003): On the putative physiological role of allopregnanolone on GABA<sub>A</sub> receptor function. *Neuropharm* 44:49–55.
- Puia G, Santi MR, Vicini S, Pritchett DB, Purdy RH, Paul SM, et al (1990): Neurosteroids act on recombinant human GABA<sub>A</sub> receptors. *Neuron* 4:759–765.
- Purdy RH, Morrow AL, Moore PH Jr, Paul SM (1991): Stress-induced elevations of gamma-aminobutyric acid type A receptor-active steroids in the rat brain. *Proc Natl Acad Sci U S A* 88:4553–4557.
- Rasmusson AM, Lipschitz DS, Wang S, Hu S, Vojvoda D, Bremner JD, et al (2001): Increased pituitary and adrenal reactivity in premenopausal women with posttraumatic stress disorder. *Biol Psychiatry* 50:965–977 [erratum appears in *Biol Psychiatry* 2002;52:771].
- Rasmusson AM, Picciotto MR, Krishnan-Sarin S (in press): Smoking as a complex but critical covariate in neurobiological studies of PTSD. A review. *J Psychopharmacol*
- Rasmusson AM, Vasek J, Lipschitz DS, Vojvoda D, Mustone ME, Shi Q, et al (2004): An increased capacity for adrenal DHEA release is associated with decreased avoidance and negative mood symptoms in women with PTSD. *Neuropsychopharm* 29:1546–1557.
- Rasmusson AM, Vythilingam M, Morgan CA III (2003): The neuroendocrinology of PTSD—new directions. *CNS Spectrums* 8:651–667.
- Resick PA, Nishith P, Weaver TL, Astin MC, Feuer CA (2002): A comparison of cognitive-processing therapy with prolonged exposure and a waiting condition for the treatment of chronic posttraumatic stress disorder in female rape victims. *J Consult Clin Psychology* 70:867–879.
- Ressler KJ, Paschall G, Zhou XL, Davis M (2002): Regulation of synaptic plasticity genes during consolidation of fear conditioning. *J Neurosci* 22:7892–902.
- Rhodes ME, McCormick CM, Frye CA (2004): 3 $\alpha$ ,5 $\alpha$ -THP mediates progesterone's effects to protect against adrenalectomy-induced cell death in the dentate gyrus of female and male rats. *Pharmacol Biochem Behav* 78:505–512.
- Romeo E, Strohle A, Spalletta G, di Michele F, Hermann B, Holsboer F, et al (1998): Effects of antidepressant treatment on neuroactive steroids in major depression. *Am J Psychiatry* 155:910–913.
- Rosenkranz JA, Moore H, Grace AA (2003): The prefrontal cortex regulates lateral amygdala neuronal plasticity and responses to previously conditioned stimuli. *J Neurosci* 23:11054–11064.
- Rupprecht R (2003): Neuroactive steroids: Mechanisms of action and neuropsychopharmacological properties. *Psychoneuroendocrin* 28:139–168.
- Rupprecht R, Reul J, Trapp T, van Steensel B, Wetzel C, Damm K, et al (1993): Progesterone receptor-mediated effects of neuroactive steroids. *Neuron* 11:523–530.
- Schmidt PJ, Daly RC, Bloch M, Smith MJ, Danaceau MA, Simpson St. Clair L, et al (2005): Dehydroepiandrosterone monotherapy in midlife-onset major and minor depression. *Arch Gen Psychiatry* 62:154–162.
- Schumacher M, Coirini H, McEwen BS (1989): Regulation of high-affinity GABA<sub>A</sub> receptors in specific brain regions by ovarian hormones. *Neuroendocrin* 50:315–320.
- Semyanov A, Walker MC, Kullmann DM (2003): GABA uptake regulates cortical excitability via cell type-specific tonic inhibition. *Nat Neurosci* 6:484–490.
- Semyanov A, Walker MC, Kullmann DM, Silver RA (2004): Tonically active GABA<sub>A</sub> receptors: Modulating gain and maintaining the tone. *Trends Neurosci* 27:262–269.
- Simms LJ, Watson D, Doebbeling BN (2002): Confirmatory factor analyses of posttraumatic stress symptoms in deployed and nondeployed veterans of the Gulf War. *J Abn Psychology* 111:637–647.
- Smith SS, Gong QH, Hsu FC, Markowitz RS, French-Mullen JM, Li X (1998): GABA(A) receptor alpha4 subunit suppression prevents withdrawal properties of an endogenous steroid. *Nature* 392:926–930.
- Sondergaard HP, Hansson L-O, Theorell T (2002): Elevated blood levels of dehydroepiandrosterone sulphate vary with symptom load in posttraumatic stress disorder: Findings from a longitudinal study of refugees in Sweden. *Psychother Psychosom* 71:298–303.
- Southwick SM, Rasmusson AM, Barron J, Arnsten A (2005): Neurobiological and neurocognitive alterations in PTSD: A focus on norepinephrine, serotonin, and the HPA axis. In: Vasterling J, Brewin C, editors. *Neuropsychology of PTSD*, New York: Guilford Publications, 27–58.
- Spivak B, Maayan R, Kotler M, Mester R, Gil-Ad I, Shtaitf B, Weizman A (2000): Elevated circulatory level of GABA<sub>A</sub>-antagonistic neurosteroids in patients with combat-related post-traumatic stress disorder. *Psychol Med* 30:1227–1231.
- Stein DJ, Seedat S, van der Linden GJ, Zungu-Dirwayi N (2000): Selective serotonin reuptake inhibitors in the treatment of post-traumatic stress disorder: A meta-analysis of randomized controlled trials. *Int Clin Psychopharm* 15(suppl 2):S31–S39.
- Strous R, Maayan R, Lapidus R, Stryker R, Lustig M, Kotler M, Weizman A (2003): Dehydroepiandrosterone augmentation in the management of negative, depressive, and anxiety symptoms in schizophrenia. *Arch Gen Psychiatry* 60:133–141.
- Sun C, Sieghart W, Kapur J (2004): Distribution of alpha1, alpha4, gamma2, and delta subunits of GABA<sub>A</sub> receptors in hippocampal granule cells. *Brain Res* 1029:207–216.
- Sundstrom-Poromaa I, Smith DH, Gong QH, Sabado TN, Li X, Light A, et al (2002): Hormonally regulated alpha(4)beta(2)delta GABA(A) receptors are a target for alcohol. *Nat Neurosci* 5:721–722.
- Trauger JW, Jiang A, Stearns BA, LoGrasso PV (2002): Kinetics of allopregnanolone formation catalyzed by human 3 $\alpha$  hydroxysteroid dehydrogenase type III (AKR1C2). *Biochem* 41:13451–13459.
- Uzunov DP, Cooper TB, Costa E, Guidotti A (1996): Fluoxetine-elicited changes in brain neurosteroid content measured by negative ion mass fragmentography. *Proc Natl Acad Sci U S A* 93:12599–12604.
- Uzunova V, Sheline Y, Davis J, Rasmusson A, Uzunova D, Costa E, Guidotti A (1998): Increase in the cerebrospinal fluid content of neurosteroids in patients with unipolar major depression who are receiving fluoxetine or fluvoxamine. *Proc Nat Acad Sci* 95:3239–3244.
- Vaiva G, Thomas P, Ducrocq F, Fontaine M, Boss V, Devos P, et al (2004): Low posttrauma GABA plasma levels as a predictive factor in the development of acute posttraumatic stress disorder. *Biol Psychiatry* 55:250–254.
- Vallee M, Rivera JD, Koob GF, Purdy RH, Fitzgerald RL (2000): Quantification of neurosteroids in rat plasma and brain following swim stress and allopregnanolone administration using negative chemical ionization gas chromatography/mass spectrometry. *Anal Biochem* 287:153–166.
- Wang S, Mason J (1999): Elevations of serum T3 levels and their association with symptoms in World War II veterans with combat-related posttraumatic stress disorder: Replication of findings in Vietnam combat veterans. *Psychosomatic Med* 61:131–138.
- Wolkowitz OM, Reus VI, Keebler A, Nelson N, Friedland M, Brizendine L, Roberts E (1999): Double-blind treatment of major depression with dehydroepiandrosterone. *Am J Psychiatry* 156:646–649.
- Yehuda R (2002): Current status of cortisol findings in post-traumatic stress disorder. *Psychiatric Clin N Amer* 25:341–368.
- Young EA, Breslau N (2004): Cortisol and catecholamines in posttraumatic stress disorder: An epidemiologic community study. *Arch Gen Psychiatry* 61:394–401.